

Second Order Perturbations in the Randall-Sundrum Infinite Brane-World Model

Hideaki Kudoh ^{*} and Takahiro Tanaka [†]

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

We discuss the non-linear gravitational interactions in the Randall-Sundrum single brane model. If we naively write down the 4-dimensional effective action integrating over the fifth dimension with the aid of the decomposition with respect to eigen modes of 4-dimensional d'Alembertian, the Kaluza-Klein mode coupling seems to be ill-defined. We carefully analyze second order perturbations of the gravitational field induced on the 3-brane under the assumption of the static and axial-symmetric 5-dimensional metric. It is shown that there remains no pathological feature in the Kaluza-Klein mode coupling after the summation over all different mass modes. Furthermore, the leading Kaluza-Klein corrections are shown to be sufficiently suppressed in comparison with the leading order term which is obtained by the zero mode truncation. We confirm that the 4-dimensional Einstein gravity is approximately recovered on the 3-brane up to second order perturbations.

PACS 04.50.+h; 98.80.Cq YITP-01-26

I. INTRODUCTION

The possibility that our 4-dimensional universe is embedded as a brane in a higher dimensional spacetime has been extensively discussed recently as the brane world scenario [1–4]. In particular, Randall and Sundrum introduced attractive models whose background bulk spacetime is 5-dimensional anti-de Sitter (AdS_5) spacetime. These models suggest the possibility of the existence of extra dimensions in the non-trivial form and the possible explanation of the large hierarchy between the Planck scale and the electroweak scale [3,4].

The behavior of the gravity in the Randall-Sundrum (RS) models has been investigated by many authors [5–14]. For the RS single brane model, which is the model with the positive tension brane alone, the extension of extra dimension is infinite. Nevertheless, the gravity on the 3-brane generated by the matter field, which is confined on the 3-brane, approximately coincides with the 4-dimensional Einstein gravity [5–8].

For the RS two-brane model whose bulk space is bounded by the positive and the negative tension branes, the 4-dimensional Einstein gravity can be recovered on both branes under the approximation of the zero mode truncation [9] when we take the stabilization mechanism of the distance between the two branes into consideration [15–17].

Although a large number of studies have been made on linear perturbations of the metric [18–23], little is known about the non-linear or non-perturbative feature of the gravity [24–30]. There is a worry about the RS single brane model. In this model, it seems that the non-linear gravitational interaction between Kaluza-Klein (KK) modes becomes strong and it diverges as we move far away from the brane. A more precise argument is that the 4-dimensional effective action including interaction terms diverges when we attempt to write it down by integrating over the dependence on the 5th direction by using the decomposition of metric perturbations in terms of eigen modes of 4-dimensional d'Alembertian [4]. This fact indicates that we cannot construct a 4-dimensional effective action for this model in an usual sense.

The above discussion is based on the analysis using the decomposition of the mass eigen modes, which is referred to as the mode-by-mode analysis. However, it was demonstrated that the mode-by-mode analysis is insufficient to deal with metric perturbations and it is necessary to take into account the contributions from all the KK modes when we discuss the regularity of linear metric perturbations at a point far from the 3-brane [11]. Thus the pathological behavior in the non-linear interaction can be expected to be an artifact solely due to the mode-by-mode analysis, although it has not been proved yet. Even if the pathological behavior might be fictitious, there is another question if the 4-dimensional Einstein gravity is recovered when we proceed to higher order perturbations?

The purpose of this paper is to give a partial answer to the above two questions. To study the non-linear behavior of the gravity, we investigate second order perturbations in the context of the RS single brane model. To simplify the analysis, we consider the static and axisymmetric configuration, which means that the 3-brane metric is spherically symmetric. Following the method developed by Garriga and Tanaka [5], we confirm that there is no pathological

^{*}E-mail: kudoh@yukawa.kyoto-u.ac.jp

[†]E-mail: tanaka@yukawa.kyoto-u.ac.jp

feature in the KK mode coupling if we sum up all the mass eigen modes, and that the contribution due to the KK mode coupling is sufficiently small compared with the leading order terms, which are obtained by the zero mode truncation. Furthermore, the result obtained by the zero mode truncation exactly agrees with the one predicted by the 4-dimensional Einstein gravity.¹

This paper is organized as follows. In the next section we first derive the second order perturbation equations in the 5-dimensional bulk, and discuss the gauge transformation as well as the boundary condition. In Sec. III, after giving brief summary on the behavior of the 5-dimensional Green function, which is need to solve the perturbation equations, we review the results for linear perturbations to give their explicit expressions in the notation of the present paper. In Sec. IV, we analyze second order perturbations of the metric induced on the 3-brane. First it is shown that the 4-dimensional Einstein gravity is recovered in the approximation of the zero mode truncation, and then we prove that the remaining contribution due to the Kaluza-Klein mode coupling can be neglected. Section V is devoted to summary.

II. PERTURBATION EQUATIONS IN THE RS MODEL

The brane world model proposed by Randall and Sundrum is composed of the 5-dimensional AdS space

$$ds^2 = g_{ab}dx^a dx^b = a^2(y)\eta_{\mu\nu}dx^\mu dx^\nu + dy^2, \quad (2.1)$$

with a single positive tension ($\sigma > 0$) 3-brane located at $y = 0$. Here $a(y) \equiv e^{-|y|/\ell}$ is the warp factor, and ℓ is the curvature radius of AdS_5 . We have denoted the 4-dimensional Minkowski metric as $\eta_{\mu\nu}$. The cosmological constant on the bulk and the tension of the 3-brane are, respectively, related to the curvature radius ℓ as $\Lambda = -6\ell^{-2}$ and as $\sigma = 3/4\pi\ell G_5$, where G_5 is 5-dimensional Newton's constant. The relation between the 5-dimensional and the 4-dimensional Newton's constants is given by $G_5 = \ell G_4$. We also use the notation $\kappa_5 \equiv 8\pi G_5$. Ordinary matter field is supposed to be localized on the brane.

In this paper, we investigate second order perturbations of this model induced by non-relativistic matter on the brane whose energy momentum tensor is given by the perfect fluid form

$$T_\mu^\nu = \text{diag}\{-\rho, P, P, P\}. \quad (2.2)$$

To simplify the analysis, we restrict our consideration to the static and axisymmetric spacetime whose axis of symmetry lies along y -direction. Namely, the 3-brane metric is spherically symmetric.

We denote the perturbed metric by $\tilde{g}_{ab} = g_{ab} + h_{ab}$. At the level of linear perturbations, it is advantageous to use the Randall-Sundrum (RS) gauge defined by

$$h_{yy} = h_{\mu y} = 0, \quad (2.3)$$

$$h^\nu_{\mu;\nu} = 0, \quad h^\mu_\mu = 0, \quad (2.4)$$

because the linear perturbation equations in this gauge take the simple form [5]

$$\mathcal{L} h_{\mu\nu} \equiv \left[a^{-2}\square^{(4)} + \partial_y^2 - 4\ell^{-2} \right] h_{\mu\nu} = 0, \quad (2.5)$$

and all components are decoupled. However, when we consider second order perturbations, we cannot impose the transverse-traceless condition (2.4) in addition to (2.3). Hence, we need to abandon to impose the condition (2.4) in second order perturbations. As a consequence, the second order perturbation equations are inevitably coupled.

¹ After we had submitted the previous version of this paper, we had the existence of Ref. [34] pointed out. In this reference, the second order metric perturbation in the region far from the star was investigated using the truncation of the first order metric perturbation at the leading order for the expansion with respect to the distance from the star. Since the Einstein equation was not solved in the whole region of the spacetime in their treatment, there remains an ambiguity to add some metric perturbations which satisfy the homogeneous linear perturbation equation only in the asymptotic region. Nevertheless, one may be able to prove that the dominant part of this remaining ambiguity can be absorbed by the redefinition of the mass parameter, and actually our results show that this is the case. Hence, the recovery of the 4-d Einstein gravity in the region far from the star is to be credited to Ref. [34]. On the other hand, what we show in the present paper is the recovery of the 4-d Einstein gravity throughout the whole spacetime including the inside of the star without assuming a specific radial matter distribution.

Here, instead of requiring (2.4), we start with the assumption of the diagonal form of metric

$$ds^2 = a^2 \left[-e^{\mathcal{A}(r,y)} dt^2 + e^{\mathcal{B}(r,y)} dr^2 + e^{\mathcal{C}(r,y)} r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \right] + dy^2, \quad (2.6)$$

which is manifestly compatible with the gauge condition (2.3), and does not loose the generality under the restriction to the static and axial-symmetric case. Furthermore, we can expect that this assumption is also compatible with the condition (2.4) at the linear order according to the result obtained in Ref. [5]. For the assumed metric form (2.6), the conditions (2.4) at the linear level, respectively, become

$$\mathcal{A}^{(1)} + \mathcal{B}^{(1)} + 2\mathcal{C}^{(1)} = 0, \quad (2.7)$$

and

$$\partial_r(r^2 \mathcal{B}^{(1)}) - 2r\mathcal{C}^{(1)} = 0, \quad (2.8)$$

where we have expanded $\mathcal{A}, \mathcal{B}$ and $\mathcal{C}$ to the second order like

$$\mathcal{A}(r, y) = \sum_{J=1,2} \mathcal{A}^{(J)}(r, y). \quad (2.9)$$

Hereafter, we neglect higher order terms without mentioning it.

Before we start to solve the 5-dimensional Einstein equation, we would like to mention the boundary condition at $y \rightarrow \infty$. An important point which we must mention here is that we are to find a solution which is regular at $y \rightarrow \infty$. If we allow the violation of regularity at the infinity, the dynamics of the RS brane world model is not uniquely determined. Then, such a model cannot be a candidate of the model that describes our universe. Hence, we require that the metric converges to the AdS_5 asymptotically. To guarantee this asymptotic condition, we simply require that the metric functions $\mathcal{A}, \mathcal{B}$ and $\mathcal{C}$ to go to 0 at $y \rightarrow \infty$. With this choice, the residual gauge degrees of freedom are completely fixed. Under this constraint, it is still possible to extend the coordinates throughout the bulk maintaining the diagonal form of the metric. We extensively refer to this gauge choice as the RS gauge because we see later that the first order quantities $\mathcal{A}^{(1)}, \mathcal{B}^{(1)}$ and $\mathcal{C}^{(1)}$ in this gauge satisfy the both conditions (2.7) and (2.8) (, although the quantities at the second order do not satisfy the transverse-traceless condition (2.4)).

A. 5D Einstein Equations in the bulk

We consider the 5-dimensional Einstein equations and derive the perturbation equations up to the second order. Since the trace of metric perturbations vanishes in the RS gauge at the linear order, it would be convenient to introduce

$$2\psi^{(J)} \equiv \mathcal{A}^{(J)} + \mathcal{B}^{(J)} + 2\mathcal{C}^{(J)}. \quad (2.10)$$

By using this quantity, the trace of metric perturbations is expressed as

$$\tilde{g}^{\mu\nu} h_{\mu\nu} = 2 \sum_{J=1,2} \psi^{(J)} - \frac{1}{2} \left[(\mathcal{A}^{(1)})^2 + (\mathcal{B}^{(1)})^2 + 2(\mathcal{C}^{(1)})^2 \right]. \quad (2.11)$$

Hence the traceless condition at the first order is simply given by $\psi^{(1)} = 0$.

The 5-dimensional vacuum Einstein equation with the cosmological term is equivalent to the following set of equations for the Ricci tensor,

$$R_y^y + \frac{4}{\ell^2} = \sum_{J=1,2} \left(\frac{2}{\ell} \psi_{,y}^{(J)} - \psi_{,yy}^{(J)} \right) - Q_{yy} = 0, \quad (2.12)$$

$$R_t^t + \frac{4}{\ell^2} = \frac{1}{2\ell} \sum_{J=1,2} \left(2\psi_{,y}^{(J)} + 4\mathcal{A}_{,y}^{(J)} - \ell \mathcal{A}_{,yy}^{(J)} - \ell a^{-2} \Delta \mathcal{A}^{(J)} \right) - \frac{1}{2a^2} S = 0, \quad (2.13)$$

$$R_r^r = \frac{1}{2r} \sum_{J=1,2} \left[\frac{1}{r^2} (r^3 \mathcal{B}^{(J)})_{,ry} - 2(r\psi_{,y}^{(J)})_{,r} + \mathcal{A}_{,y}^{(J)} \right] + \frac{1}{2} \psi_{,r}^{(1)} \mathcal{B}_{,y}^{(1)} - Q_{ry} = 0, \quad (2.14)$$

where $\Delta \equiv \sum_{i=1}^3 \partial_i^2$ and we have defined

$$S(r, y) \equiv -\frac{1}{r^2} \partial_r \left(r^2 \mathcal{A}_{,r}^{(1)} \mathcal{B}^{(1)} \right) + a^2 \psi_{,y}^{(1)} \mathcal{A}_{,y}^{(1)} + \psi_{,r}^{(1)} \mathcal{A}_{,r}^{(1)}, \quad (2.15)$$

$$Q_{\mu\nu}(r, y) \equiv \frac{1}{4} \left(\mathcal{A}_{,\mu}^{(1)} \mathcal{A}_{,\nu}^{(1)} + \mathcal{B}_{,\mu}^{(1)} \mathcal{B}_{,\nu}^{(1)} + 2\mathcal{C}_{,\mu}^{(1)} \mathcal{C}_{,\nu}^{(1)} \right). \quad (2.16)$$

Note that no other equations for the remaining components are independent.

First we consider the trace part $\psi^{(J)}$, which can be evaluated by integrating (2.12). From the requirement of the boundary condition, $\psi^{(J)}$ must go to 0 at $y \rightarrow \infty$. Thus, we obtain

$$\psi^{(J)}(r, y) = -\epsilon^{(J)} \int_{\infty}^y dy'' e^{2y''/\ell} \int_{\infty}^{y''} dy' e^{-2y'/\ell} Q_{yy}. \quad (2.17)$$

where we have introduced a symbol $\epsilon^{(J)}$ that is defined by $\epsilon^{(1)} = 0$ and $\epsilon^{(2)} = 1$ to represent the first and the second order equations in a single expression. As is anticipated above, we can see from Eq. (2.17) that the traceless condition at the linear order is actually satisfied, while that at the second order cannot be imposed any longer in general.

Let us now turn to $\mathcal{B}^{(J)}$. Integrating (2.14), we obtain

$$\mathcal{B}^{(J)}(r, y) = -\frac{1}{r^3} \int r^2 \mathcal{A}^{(J)} dr + \frac{2}{r^3} \epsilon^{(J)} \left(\int dr r^2 (r\psi^{(2)})_{,r} + \int_{\infty}^y dy \int dr r^3 Q_{ry} \right). \quad (2.18)$$

As before, the integration constant is fixed by the boundary condition at $y \rightarrow \infty$. Now, combined with Eq. (2.7), it is easy to see that Eq.(2.8) holds at the linear order. Hence, it is confirmed that our choice of gauge is equivalent to the RS gauge.

So far, we obtained the relations between the metric functions $\mathcal{A}^{(J)}$, $\mathcal{B}^{(J)}$ and $\mathcal{C}^{(J)}$. Substituting these into Eq. (2.13), we obtain a single equation for $\mathcal{A}^{(J)}$,

$$\mathcal{L}[a^2 \mathcal{A}^{(J)}] = \epsilon^{(J)} [2a^2 \ell^{-1} \psi_{,y} - S]. \quad (2.19)$$

The source terms are absent at the linear order, as they are composed of the quadratic in the linear order quantities. Since $\psi^{(1)} = 0$, S defined in Eq.(2.15) simplifies as

$$S(r, y) = \frac{1}{r^2} \partial_r \left(r^2 \mathcal{A}_{,r}^{(1)} \mathcal{B}^{(1)} \right). \quad (2.20)$$

B. Boundary condition

In the previous subsection, we have obtained the master equation for the metric functions in the bulk up to the second order. To solve this equation, we need to know the boundary condition to be imposed on the 3-brane. The boundary condition on the 3-brane is specified by the Israel's junction condition [31]. However, in the RS gauge defined above, the y -constant surface is, by construction, chosen so that the metric functions go to 0 at $y \rightarrow \infty$. After fixing the y -constant surface for large y , coordinates are extended to the region near the 3-brane. Therefore, we can no longer expect that the location of the 3-brane coincides with the $y = 0$ surface in general [5]. In such coordinates, the junction condition is not so trivial. Thus it will be convenient to introduce other coordinates $\bar{x}^a$, in which the location of the 3-brane stays at $\bar{y} = 0$ but the metric form is still kept diagonal. We associate a bar with the quantities written in these coordinates, say, $\bar{\mathcal{A}}$. We denote this choice of coordinates by the Gaussian normal (GN) gauge. By construction, $\bar{h}_{\mu\nu}|_{\bar{y}=0}$ gives the 4-metric induced on the 3-brane.

The junction condition on the 3-brane is simply written in the GN gauge as ²

$$\tilde{g}^{\gamma\nu}(\bar{x}) \partial_{\bar{y}} (a^{-2}(\bar{y}) \bar{h}_{\mu\nu}(\bar{x})) = -\kappa_5 \left(T^{\gamma}_{\mu} - \frac{1}{3} \delta^{\gamma}_{\mu} T^{\lambda}_{\lambda} \right) (\bar{x}). \quad (\text{at } \bar{y} = 0+),$$

because this is a relation for the quantities in the GN gauge. However, $f(x)|_{y=0} = 0$ is obviously equivalent to $f(\bar{x})|_{\bar{y}=0} = 0$ for any function f . To stress this point, we always use the coordinates without bar hereafter. The only exception is Eq. (2.24).

$$\tilde{g}^{\gamma\nu}(x)\partial_y(a^{-2}(y)\bar{h}_{\mu\nu}(x)) = -\kappa_5 \left(T^\gamma{}_\mu - \frac{1}{3}\delta^\gamma{}_\mu T^\lambda{}_\lambda \right) (\mathbf{x}). \quad (\text{at } y=0+) \quad (2.21)$$

As mentioned earlier, we assume the energy momentum tensor of the perfect fluid form (2.2). Then, the 4-dimensional energy-momentum conservation $T^\nu{}_{\mu;\nu} = 0$ becomes

$$(\rho(r) + P(r))\partial_r \bar{\mathcal{A}}^{(1)}(r, 0) + 2\partial_r P(r) = 0, \quad (2.22)$$

and hence we find that $P(r)$ is a second order quantity. This equation represents the force balance between pressure and gravity acting on the matter field.

Taking it into account that $P(r)$ is second order and expanding ρ as $\rho = \rho^{(1)} + \rho^{(2)}$, the explicit junction conditions for the metric functions become

$$\begin{aligned} \partial_y \bar{\mathcal{A}}^{(J)} &= \kappa_5 \left(\frac{2}{3}\rho^{(J)} + \epsilon^{(J)}P \right), \\ \partial_y \bar{\mathcal{B}}^{(J)} &= \partial_y \bar{\mathcal{C}}^{(J)} = -\frac{\kappa_5}{3}\rho^{(J)}, \quad (\text{at } y=0+). \end{aligned} \quad (2.23)$$

The boundary condition obtained above is written in terms of the variables in the GN gauge. To interpret the conditions (2.23) in terms of the variables in the RS gauge, we consider the gauge transformation between these two gauges, which is defined by

$$\tilde{g}_{ab}(\bar{x}) d\bar{x}^a d\bar{x}^b = \tilde{g}_{ab}(x) dx^a dx^b, \quad (2.24)$$

with $\bar{x}^a = x^a + \xi^a(x)$. Since $h_{y\mu}$ and h_{yy} vanish in both gauges, the infinitesimal gauge transformation $\xi^a = \xi^{(1)a} + \xi^{(2)a}$ between them is restricted to the following form

$$\xi^y(r, y) = \hat{\xi}^y(r) - \epsilon^{(J)} \left[\frac{\ell}{4a^2} (\hat{\xi}_{,r}^y)^2 \right], \quad (2.25)$$

$$\xi^r(r, y) = \hat{\xi}^r(r) - \frac{\ell}{2a^2} \hat{\xi}_{,r}^{(J)} + \epsilon^{(J)} \hat{\xi}_{,r}^{(1)} \left[\frac{\ell}{2a^2} (\hat{\xi}_{,r}^{(1)} - \frac{2}{\ell} \hat{\xi}^y) + \int_0^y \frac{dy'}{a^2} \bar{\mathcal{B}}^{(1)}(r, y') \right]. \quad (2.26)$$

Functions $\hat{\xi}^y(r)$ and $\hat{\xi}^r(r)$ are integration constants which appear as a result of y -integration. We discuss how these functions are determined soon later.

We denote the difference between the metric in the RS gauge and that in the GN gauge as

$$\delta\mathcal{A}^{(J)}(r, y) \equiv \mathcal{A}^{(J)}(r, y) - \bar{\mathcal{A}}^{(J)}(r, y). \quad (2.27)$$

The remaining metric functions $\delta\mathcal{B}^{(J)}$ and $\delta\mathcal{C}^{(J)}$ are defined in the same way. Then, the gauge transformations are given by

$$\begin{aligned} \delta\mathcal{A}^{(J)} &= -\frac{2}{\ell} \hat{\xi}^{(J)} + \epsilon^{(J)} \left[\bar{\mathcal{A}}_{,y}^{(1)} \hat{\xi}^{(1)} + \bar{\mathcal{A}}_{,r}^{(1)} \hat{\xi}^r \right], \\ \delta\mathcal{B}^{(J)} &= -\frac{2}{\ell} \hat{\xi}^y + 2\hat{\xi}_{,r}^{(J)} + \epsilon^{(J)} \left[\bar{\mathcal{B}}_{,y}^{(1)} \hat{\xi}^{(1)} + \bar{\mathcal{B}}_{,r}^{(1)} \hat{\xi}^r - \left(\hat{\xi}_{,r}^{(1)} \right)^2 + e^{2y/\ell} \left(\hat{\xi}_{,r}^{(1)} \right)^2 \right], \\ \delta\mathcal{C}^{(J)} &= -\frac{2}{\ell} \hat{\xi}^y + \frac{2}{r} \hat{\xi}^r + \epsilon^{(J)} \left[\bar{\mathcal{C}}_{,y}^{(1)} \hat{\xi}^{(1)} + \bar{\mathcal{C}}_{,r}^{(1)} \hat{\xi}^r - \left(\frac{1}{r} \hat{\xi}^r \right)^2 \right]. \end{aligned} \quad (2.28)$$

Now we are ready to derive the equation that determines $\hat{\xi}^y(r)$. We evaluate the identity $\psi_{,y} = \bar{\psi}_{,y} + \delta\psi_{,y}$ at $y=0$. The expression for the left hand side is obtained by means of Eq.(2.17), while the right hand side is evaluated by substituting Eqs. (2.23), (2.25), (2.26) and (2.28). After tedious but straightforward computation, we obtain

$$\Delta \hat{\xi}^y = \frac{\kappa_5}{6} T^{(J)} + \epsilon^{(J)} \left[\Xi(r) - \psi_{,y}|_{y=0} \right], \quad (2.29)$$

with

$$\Xi(r) \equiv \Delta \left[\frac{1}{\ell} (\hat{\xi}^y)^2 + \int_\infty^r \hat{\xi}_{,r'}^{(1)} \mathcal{B}^{(1)}(r', 0) dr' \right] - \frac{3}{\ell} (\hat{\xi}_{,r}^{(1)})^2 + \hat{\xi}^r \partial_r \Delta \hat{\xi}^y + \frac{\ell}{2} (\hat{\xi}_{,rr}^{(1)})^2 + \frac{\ell}{r^2} (\hat{\xi}_{,r}^{(1)})^2 \quad (2.30)$$

The gauge freedom for the radial coordinate in the GN gauge has not been fixed. Although the simplest choice might be to take $\xi^{(J)r}$ to vanish on the brane, for later convenience we impose the isotropic gauge condition on the 3-brane, $\bar{\mathcal{B}} = \bar{\mathcal{C}}$ at $\bar{y} = 0$. Rewriting this condition by using $\mathcal{B}$ and $\mathcal{C}$ with the substitution of (2.28), we obtain

$$\xi^{(1)r} = -\frac{r}{4}\mathcal{B}^{(1)}, \quad (2.31)$$

where we have used (2.8). Here we show the result only for the first order, because we do not need the explicit expression for $\xi^{(2)r}$ in the following discussion.

Once we obtain an explicit expression for $\hat{\xi}^{(J)y}$, it is easy to derive the boundary condition for the metric functions in the RS gauge from the junction condition in the GN gauge (2.23). Especially, the boundary condition required to solve the master equation (2.19) is deduced by substituting the relations obtained above into the right hand side of the equation, $\mathcal{A}_{,y} = \bar{\mathcal{A}}_{,y} + \delta\mathcal{A}_{,y}$. Imposing this boundary condition is equivalent to add a δ -function source localized on the 3-brane. The 5-dimensional master equation (2.19) including the boundary condition becomes

$$[\mathcal{L} + 4\ell^{-1}\delta(y)](a^2\mathcal{A}^{(J)}) = 2\kappa_5 \Sigma^{(J)}(r)\delta(y) + \epsilon^{(J)}[2a^2\ell^{-1}\psi_{,y} - S], \quad (2.32)$$

where

$$\begin{aligned} \Sigma^{(J)}(r) &\equiv \kappa_5^{-1}\partial_y\mathcal{A}^{(J)}|_{y=0} \\ &= \frac{2}{3}\rho^{(J)} + \epsilon^{(J)}\left[P + \kappa_5^{-1}\partial_y(\delta\mathcal{A}^{(2)})|_{y=0}\right], \end{aligned} \quad (2.33)$$

By using the Green function which satisfies

$$[\mathcal{L} + 4\ell^{-1}\delta(y)]G(\mathbf{x}, y; \mathbf{x}', y') = \delta(y - y')\delta^3(\mathbf{x} - \mathbf{x}'), \quad (2.34)$$

the formal solution for the master equation (2.32) is given by

$$\begin{aligned} a^2\mathcal{A}^{(J)}(r, y) &= 2\kappa_5 \int d^3x' G(\mathbf{x}, y; \mathbf{x}', 0) \Sigma^{(J)} \\ &\quad - 2\epsilon^{(J)} \int d^3x' \int_0^\infty dy' G(\mathbf{x}, y; \mathbf{x}', y') \left[S(r', y') + \frac{2}{\ell} \int_\infty^{y'} a^2(y'') Q_{yy}(r', y'') dy'' \right]. \end{aligned} \quad (2.35)$$

The factor 2 in the second term of (2.35) comes from Z_2 -symmetry. In §III we discuss some basic aspects of the Green function, and solve the above equation at the linear order. In §IV we extend our analysis to the second order.

III. GREEN FUNCTION AND FIRST ORDER PERTURBATIONS

A. Green function

We need the Green function to evaluate explicitly the formal solution (2.35). In the static case Green function is

$$G(\mathbf{x}, y; \mathbf{x}', y') = -\frac{1}{4\pi\ell R} \left(a^2(y)a^2(y') + \ell \int_0^\infty u_m(y)u_m(y')e^{-mR}dm \right), \quad (3.1)$$

where $u_m(y)$ is the mode function, and we have introduced the notation

$$R \equiv |\mathbf{x} - \mathbf{x}'|.$$

The explicit form of the mode function is given in terms of Bessel functions as

$$u_m(y) = N_m [J_1(m\ell)Y_2(m\ell/a) - Y_1(m\ell)J_2(m\ell/a)], \quad (3.2)$$

with $N_m = \sqrt{m\ell}/\sqrt{2(J_1(m\ell)^2 + Y_1(m\ell)^2)}$. It is orthonormalized as

$$2 \int_0^\infty \frac{dy}{a^2} u_m u_{m'} = \delta(m - m'). \quad (3.3)$$

In particular, setting $m' = 0$, we have

$$\int_0^\infty dy u_m = 0 \quad (m \neq 0). \quad (3.4)$$

The first term on the right hand side of (3.1) is the contribution from the zero mode whose 4-dimensional mass eigenvalue is zero ($m = 0$). We denote this part of Green function by G_0 . The second term corresponds to the propagator due to the Kaluza-Klein states which have nonzero mass eigenvalues ($m > 0$). This term is denoted by G_K . Thus, $G = G_0 + G_K$.

For large separation $R \gg \ell$, the existence of the factor e^{-mR} in the second term in (3.1), i.e. G_K , implies that the integral over m is dominated by the contribution from small m . When the source is on the 3-brane ($y' = 0$), we can approximately evaluate G_K by expanding Bessel functions taking $m\ell$ as small, but here the Bessel function with the argument $m\ell/a$ is to be kept unexpanded because a can be exponentially small. Then, the integration over m can be performed for the leading power of m explicitly. The result is given by [5]

$$G(\mathbf{x}, y; \mathbf{x}', 0) \approx -\frac{a^3}{8\pi\ell} \frac{(2a^2 R^2 + 3\ell^2)}{(a^2 R^2 + \ell^2)^{3/2}}. \quad (3.5)$$

For a small separation $R \ll \ell$, the Green function is dominated by the contribution from modes with large m . In this limit, the Green function behaves as the ordinary 5-dimensional one $\approx -1/4\pi^2 [R^2 + \ell^2(1 - a^{-1})^2]$.

When we discuss second order perturbations later, we use the following inequalities

$$0 \leq -G(\mathbf{x}, y; \mathbf{x}', 0) \leq \frac{a^2(y)}{4\pi\ell R^2} (R + \ell), \quad (3.6)$$

This inequality is suggested by the asymptotic form of the Green function, and it is confirmed by numerical calculations as shown in Fig.1.




FIG. 1. We calculated numerically G and $a^2(R + \ell)/(4\pi\ell R^2) + G$ to show the inequalities of (3.6). Since there appears no logarithmic divergence, G and $a^2(R + \ell)/(4\pi\ell R^2) + G$ are positive definite at least in the region shown in the figures.

B. linear perturbations

We start with linear perturbations. From Eq.(2.35), $\mathcal{A}^{(1)}$ is given by

$$a^2 \mathcal{A}^{(1)} = \frac{4\kappa_5}{3} \int d^3 x' G(\mathbf{x}, y; \mathbf{x}', 0) \rho^{(1)}(r'). \quad (3.7)$$

Suppose that r_* is the radius of the star. When we consider the metric at a point outside the star $r - r_* \gg \ell$ or far from the 3-brane, we can safely replace G by the approximation (3.5) with the relative error of $O(\ell^2/r_*^2)$ ³. Furthermore, at a field point far from the star, the matter distribution can be replaced with a point source. Then, we obtain $\mathcal{A}^{(1)}(r, y) \sim -4 G_4 M a (2a^2 r^2 + 3\ell^2) / 3(a^2 r^2 + \ell^2)^{3/2}$. By using Eq. (2.18), we find $\mathcal{B}^{(1)}(r, y) \sim 4 G_4 M a / 3 \sqrt{a^2 r^2 + \ell^2}$.

On the other hand, if we are interested in the metric induced on the brane, we can also set $y = 0$. Even if we consider the interior of the star, the inequality (3.6) implies that the contribution to $\mathcal{A}^{(1)}$ from G_K is, at most, of $O((\kappa_4 M_*/r_*)(\ell^2/r_*^2) \log(\ell/r_*))$, where $M_* \equiv 4\pi \int r^2 \rho(r) dr$. Hence, it is small by a factor of $O((\ell^2/r_*^2) \log(\ell/r_*))$ compared with the leading term. Then, neglecting the collections of this order or higher, we obtain

$$\mathcal{A}^{(1)}(r, 0) \approx \frac{8}{3} \phi, \quad (3.8)$$

$$\mathcal{B}^{(1)}(r, 0) \approx -\frac{8}{3r} \partial_r \Delta^{-1} \phi, \quad (3.9)$$

where Δ^{-1} is the inverse of the Laplacian operator, and we have introduced Newtonian potential

$$\phi \equiv 4\pi G_4 \Delta^{-1} \rho. \quad (3.10)$$

These are transformed to the isotropic GN gauge by using (2.28) with

$$\hat{\xi}^{(1)y} = -\frac{\ell \phi}{3}, \quad \hat{\xi}^{(1)r} \approx \frac{2}{3} \partial_r \Delta^{-1} \phi, \quad (3.11)$$

which are derived from (2.29) and (2.31), respectively. Finally, the resulting metric functions turn out to be

$$-\bar{\mathcal{A}}^{(1)}(r, 0) \approx \bar{\mathcal{B}}^{(1)}(r, 0) = \bar{\mathcal{C}}^{(1)}(r, 0) \approx -2\phi(r), \quad (3.12)$$

which agree with the result for the 4-dimensional Einstein gravity.

IV. SECOND ORDER PERTURBATIONS

For later convenience, we quote the result obtained in the preceding section as

$$\mathcal{A}^{(2)}(r, 0) \equiv A_\Sigma + A_S + A_Q - \frac{2}{\ell} \Delta^{-1} \int_0^\infty dy' a^2 Q_{yy}, \quad (4.1)$$

where

$$\begin{aligned} A_\Sigma &= 2\kappa_5 \int d\mathbf{x}'^3 G(\mathbf{x}, 0; \mathbf{x}', 0) \Sigma^{(2)}, \\ A_S &= 2 \int d^3 x' \int_0^\infty dy' G(\mathbf{x}, 0; \mathbf{x}', y') \frac{1}{r'^2} \partial_{r'} \left(r'^2 \mathcal{A}_{,r'}^{(1)}(r', y') \mathcal{B}^{(1)}(r', y') \right), \\ A_Q &= \frac{4}{\ell} \int d^3 x' \int_0^\infty dy' \left[\int_\infty^{y'} G(\mathbf{x}, 0; \mathbf{x}', y'') dy'' \right] a^2(y') Q_{yy}(r', y'). \end{aligned} \quad (4.2)$$

The source term $\Sigma^{(2)}$ for A_Σ is given in Eq.(2.33). The third term in (2.35), is separated into two pieces, A_Q and the last term in (4.1), by performing integration by parts. To obtain the expression of this last term, we have also used $\int d^3 x' \int_0^\infty G dy' = \int d^3 x' \int_0^\infty G_0 dy' = (1/2) \Delta^{-1}$. Here the first equality follows from Eq. (3.4).

Since we are interested in the metric induced on the 3-brane, we also write down the expression for $\bar{\mathcal{A}}^{(2)}$ at $y = 0$. Combining (2.25), (2.28), (2.29) and (4.1), we obtain

$$\bar{\mathcal{A}}^{(2)}(r, 0) = A_\Sigma + A_S + A_Q + \frac{\kappa_4}{3} \Delta^{-1} T^{(2)} + \left[\frac{2}{\ell} \Delta^{-1} \Xi - \frac{1}{2} \left(\hat{\xi}_{,r}^{(1)y} \right)^2 - \bar{\mathcal{A}}_{,y}^{(1)} \hat{\xi}^{(1)y} - \bar{\mathcal{A}}_{,r}^{(1)} \hat{\xi}^{(1)r} \right]_{y=0}. \quad (4.3)$$

³Here r_* in the denominator is just inserted to adjust the dimensionality. It can be r instead of r_* .

Here we note that the last term in (4.1) is canceled with a term which arises from the gauge transformation.

The terms in the square brackets are evaluated just by substituting the estimate at the first order, and the first term A_Σ has structure similar to first order perturbations, i.e., the source term is localized on the brane. Thus, the evaluation of these terms is straightforward. What needs detailed investigation is the evaluation of A_S and A_Q . Since the source terms of A_S and A_Q , which are quadratic in the linear perturbation quantities, distribute through the 5-dimensional bulk, it is necessary to evaluate a convolution of three 5-dimensional Green functions.

Deferring the estimate of this convolution until § IV B, let us turn to the spatial components of the second order metric perturbations. Although each spatial component depends on the choice of $\xi^{(2)r}$, the gauge invariant combination

$$\begin{aligned} \left\{ \bar{\mathcal{B}}^{(2)} - \partial_r(r\bar{\mathcal{C}}^{(2)}) \right\}_{y=0} &= \left\{ \mathcal{B}^{(2)} - \partial_r(r\mathcal{C}^{(2)}) - \delta\mathcal{B}^{(2)} + \partial_r(r\delta\mathcal{C}^{(2)}) \right\}_{y=0} \\ &= \left\{ \frac{r}{2}\bar{\mathcal{A}}_{,r}^{(2)} + \frac{r}{2}\partial_r\delta\mathcal{A}^{(2)} - \delta\mathcal{B}^{(2)} + \partial_r(r\delta\mathcal{C}^{(2)}) \right\}_{y=0} - r \int_0^\infty dy Q_{yr} \end{aligned} \quad (4.4)$$

does not contain $\xi^{(2)r}$. Here we have used Eqs. (2.10) and (2.18) with $y = 0$ to eliminate $\mathcal{B}$ and $\mathcal{C}$. Taking the isotropic gauge $\bar{\mathcal{B}}^{(2)} = \bar{\mathcal{C}}^{(2)}$, the left hand side becomes $-r\bar{\mathcal{B}}_{,r}^{(J)}$. Integrating this equation with respect to r , we obtain the expression for $\bar{\mathcal{B}}^{(2)} (= \bar{\mathcal{C}}^{(2)})$. Here we note that the terms in the square brackets in Eq.(4.4) contains $\xi^{(2)y}$, from which a term with y -integration,

$$-\frac{3}{\ell} \int_0^\infty dy a^2 Q_{yy},$$

arises. Combined with the last term with y -integration in Eq. (4.4), this term is reduced to the expression that does not contain y -integration. The detail of calculation is explained in Appendix A. Here we just quote the final result

$$\bar{\mathcal{B}}^{(2)}(r, 0) = -\frac{1}{2}\bar{\mathcal{A}}^{(2)} + \Delta^{-1} \left(\frac{\kappa_4}{2}T^{(2)} + \frac{3}{\ell}\Xi + \frac{1}{2}Q_{yy} - \frac{3}{16}(\mathcal{A}_{,r}^{(1)})^2 \right) + B_\Sigma, \quad (\text{at } y = 0), \quad (4.5)$$

where

$$B_\Sigma \equiv -\frac{3}{4}(\hat{\xi}_{,r}^{(1)y})^2 + \int dr \frac{1}{r}(\hat{\xi}_{,r}^{(1)y})^2 + \frac{\ell\kappa_4}{4}\Delta^{-1} \left[(3\mathcal{A}_{,y}^{(1)} + \mathcal{B}_{,y}^{(1)})\Sigma^{(1)} - \frac{1}{r^3}(3\mathcal{B}_{,y}^{(1)} + \mathcal{A}_{,y}^{(1)}) \int r^2 \Sigma^{(1)} dr \right]_{y=0}.$$

We find that the expression is reduced to a closed form written solely in terms of the quantities on the 3-brane except for $\bar{\mathcal{A}}^{(2)}(r, 0)$.

A. Recovery of the 4-dimensional Einstein gravity

In this subsection, we evaluate the metric induced on the 3-brane at the leading order in ℓ/r_* , and show that the result completely agrees with the one predicted by the 4-dimensional Einstein gravity.

As for A_Σ , since the source term is localized on the 3-brane, we can approximate it as

$$A_\Sigma \approx 2\kappa_4\Delta^{-1}\Sigma^{(2)}, \quad (4.6)$$

which is justified for the same reason explained in evaluating the induced metric at the linear order.

As mentioned earlier, the point which needs careful analysis is the computation of A_S and A_Q . To evaluate these terms we need to evaluate a convolution of the 5-dimensional Green functions. Since G is composed of G_0 and G_K , the contribution from these terms is decomposed into several pieces depending on which combination of three propagators is used. For example, there is a mode coupling in which a zero mode propagator G_0 propagates the second order source produced by a product of KK mode contributions. We denote this mode coupling as

$$[K, K; 0],$$

and similar labels are assigned for the other mode couplings, too.

Here in the present subsection, we simply neglect the pieces containing the KK propagator G_K . Namely, we just take into account $A_{S[0,0;0]}$ and $A_{Q[0,0;0]}$. The justification of this approximation is given in the next subsections,

where we prove that the neglected pieces in A_S and A_Q containing G_K are actually suppressed by a factor of $O((\ell^2/r_*^2) \log(r_*/\ell))$.

As long as only the contributions from the zero mode are concerned, the first order metric functions are all constant in y . Since the source term of A_Q contains differentiation of the first order metric functions with respect to y , we find that $A_{Q[0,0;0]} \approx 0$. As for $A_{S[0,0;0]}$, after the y -integration, we find that it is reduced to

$$A_S \approx \Delta^{-1} \frac{1}{r^2} \partial_r \left(r^2 \mathcal{A}_{,r}^{(1)} \mathcal{B}^{(1)} \right). \quad (4.7)$$

Now that the evaluation of $\bar{\mathcal{A}}^{(2)}(r, 0)$ is straightforward. Substituting the evaluations of the first order quantities presented in § III B, we finally obtain

$$\bar{\mathcal{A}}^{(2)}(r, 0) \approx \kappa_4 \Delta^{-1} \left(\rho^{(2)} - 2\phi \rho^{(1)} + 3P \right), \quad (4.8)$$

and

$$\bar{\mathcal{B}}^{(2)}(r, 0) = \bar{\mathcal{C}}^{(2)}(r, 0) \approx -\Delta^{-1} \left[\kappa_4 \left(\rho^{(2)} - 2\phi \rho^{(1)} \right) + (\phi_{,r})^2 \right]. \quad (4.9)$$

It must be noted that B_Σ in (4.5) gives only higher order correction of $O(\ell^2 \kappa_4^2)$ as it is easily shown by using (2.23), (2.28), (2.29) and (2.31). These results agree with those for the 4-dimensional Einstein gravity, whose brief derivation is given in Appendix B.

B. Suppression of the KK-mode propagation

In the preceding subsection the terms in A_S and A_Q containing a KK-mode propagator were neglected. Here we show that the contribution from these terms is in fact negligible. We begin with discussing rather general things. As for A_S , the terms $A_{S[0,0;K]}$, $A_{S[0,K;0]}$ and $A_{S[K,0;0]}$ vanish because of the orthogonality (3.4). Hence the terms to be investigated are 1.) $A_{S[K,K;0]}$, $A_{S[0,K;K]}$, $A_{S[K,0;K]}$ and $A_{S[K,K;K]}$. As for A_Q , the situation is a little simpler. Recall the fact mentioned above that the zero mode contribution in the first order metric functions is y -independent. Thus we can say that $A_{Q[0,*,*]}$ and $A_{Q[*,0,*]}$ vanish. Therefore, all the terms that we need to consider for A_Q are 2.) $A_{Q[K,K;0]}$ and $A_{Q[K,K;K]}$.

1. $A_{S[K,K;0]}$, $A_{S[K,0;K]}$ and $A_{S[0,K;K]}$

The source term of A_S is rewritten as

$$\frac{1}{r^2} \partial_r \left(r^2 \mathcal{A}_{,r}^{(1)} \mathcal{B}^{(1)} \right) = \frac{1}{r_*^2} \left[\left\{ \left(r_*^2 \partial_r^2 - \frac{r_*^2}{r} \partial_r \right) \mathcal{A}^{(1)} \right\} \left(-\frac{1}{r^3} \int^r dr r^2 \mathcal{A}^{(1)} \right) - \left(\frac{r_*^2}{r} \partial_r \mathcal{A}^{(1)} \right) \mathcal{A}^{(1)} \right]. \quad (4.10)$$

Each source term has the form of

$$\left(\mathcal{O}_r^{[1]} \mathcal{A}^{(1)} \right) \left(\mathcal{O}_r^{[2]} \mathcal{A}^{(1)} \right) \quad (4.11)$$

with $\mathcal{O}_r^{[1,2]} = 1$, $\frac{r_*^2}{r} \partial_r$, $r_*^2 \partial_r^2$ or $r^{-3} \int_0^r dr r^2$. By using the orthonormal condition (3.3), we obtain the formula

$$\int_0^\infty dy a^{-2} G_K(\mathbf{x}_1, y; \mathbf{x}_2, 0) G_K(\mathbf{x}_3, y; \mathbf{x}_4, 0) = \frac{1}{2(4\pi)^2 R_{12} R_{34}} \mu(R_{12} + R_{34}), \quad (4.12)$$

where $R_{AB} \equiv |\mathbf{x}_A - \mathbf{x}_B|$ and

$$\mu(s) \equiv \int_0^\infty dm u_m(0)^2 e^{-ms}. \quad (4.13)$$

The function $\mu(s)$ is bounded by

$$0 \leq \mu(s) < \frac{\ell}{2s(s + \ell)}. \quad (4.14)$$

We begin with $A_{S[K,K;0]} + A_{S[K,K;K]}$. We can derive an upper bound for the absolute value of each term as

$$\begin{aligned} & \left| \frac{2}{r_*^2} \int d^3x' \int_0^\infty dy' G(\mathbf{x}, 0; \mathbf{x}', y') \left[\mathcal{O}_{r_1}^{[1]} \mathcal{A}_K^{(1)}(r_1, y') \mathcal{O}_{r_3}^{[2]} \mathcal{A}_K^{(1)}(r_3, y') \right]_{r_1=r_3=r'} \right| \\ & < \frac{1}{8\pi\ell r_*^2} \int d^3x' \int_0^\infty dy' \frac{a^2(y')}{R^2} (R + \ell) \left[\left\{ \mathcal{O}_{r_1}^{[1]} \mathcal{A}_K^{(1)}(r_1, y') + \mathcal{O}_{r_1}^{[2]} \mathcal{A}_K^{(1)}(r_1, y') \right\}^2 \right. \\ & \quad \left. + \left\{ \mathcal{O}_{r_1}^{[1]} \mathcal{A}_K^{(1)}(r_1, y') - \mathcal{O}_{r_1}^{[2]} \mathcal{A}_K^{(1)}(r_1, y') \right\}^2 \right]_{r_1=r'}, \end{aligned} \quad (4.15)$$

where $\mathcal{A}_K^{(1)}$ is a part of $\mathcal{A}^{(1)}$ propagated by the KK mode propagator, and we have used (3.6). In below, we show that the each term on the right hand side of the above inequality is at most $O(\ell^2 \log(r_*/\ell))$. We express the respective term as

$$\frac{1}{r_*^2} \int d^3x' \frac{R + \ell}{R^2} \left[\mathcal{O}_{r_1}^{[1]} \mathcal{O}_{r_3}^{[2]} v(r_1, r_3) \right]_{r_1=r_3=r'}, \quad (4.16)$$

with

$$\begin{aligned} v(r_1, r_3) &\equiv \ell^{-1} \int_0^\infty dy a^2 \mathcal{A}_K^{(1)}(r_1) \mathcal{A}_K^{(1)}(r_2) \\ &= \frac{\kappa_4^2 \ell}{2r_1 r_3} \int_0^{r_*} dr_2 r_2 \Sigma^{(1)}(r_2) \int_0^{r_*} dr_4 r_4 \Sigma^{(1)}(r_4) \int_{|r_1-r_2|}^{r_1+r_2} dR_{12} \int_{|r_3-r_4|}^{r_3+r_4} dR_{34} \mu(R_{12} + R_{34}), \end{aligned} \quad (4.17)$$

where $v(r_1, r_3)$ has been rewritten by using Eq. (4.12). Introducing $U(s) \equiv -\int_0^s ds' \int_\infty^{s'} ds'' \mu(s'')$,

$$\begin{aligned} v(r_1, r_3) &= -\frac{\kappa_4^2 \ell}{2r_1 r_3} \int_0^{r_*} dr_2 r_2 \Sigma^{(1)}(r_2) \int_0^{r_*} dr_4 r_4 \Sigma^{(1)}(r_4) \left(U(r_1 + r_2 + r_3 + r_4) - U(|r_1 - r_2| + r_3 + r_4) \right. \\ & \quad \left. - U(r_1 + r_2 + |r_3 - r_4|) + U(|r_1 - r_2| + |r_3 - r_4|) \right). \end{aligned} \quad (4.18)$$

By construction, $U(0) = 0$. The inequality (4.14) indicates that

$$0 \leq U(s) \leq \frac{1}{2} \left[s \log \left(1 + \frac{\ell}{s} \right) + \ell \log \left(1 + \frac{s}{\ell} \right) \right]. \quad (4.19)$$

As long as $s \lesssim r_*$, we find that $U(s)$ is at most $O(\ell \log(r_*/\ell))$.

Let us consider the action of once or twice differentiation with respect to r_1 on U . Since r_1 appears in the arguments of U only in the combination of $r_1 + r_2$ or $|r_1 - r_2|$, the differentiation with respect to r_1 can be replaced with that to r_2 . Then, integrating by parts, finally the differentiation can be moved so that it acts on the source term $r_2 \Sigma^{(1)}(r_2)$. As long as first or second derivative is concerned, differentiation of U does not appear as the boundary term. The same thing is true for the pair of r_3 and r_4 . Now, using the bound for U obtained above, we can conclude that $\mathcal{O}_{r_1}^{[1]} \mathcal{O}_{r_3}^{[2]} v(r_1, r_3)$ is at most $O(\ell^2 \log(r_*/\ell))$ for $r_1, r_3 \lesssim r_*$. When $r_1, r_3 \gtrsim r_*$, the argument of μ cannot be small. Then, we can use the bound $\mu(s) < \ell/2s^2$. Therefore, $\mathcal{O}_{r_1}^{[1]} \mathcal{O}_{r_3}^{[2]} v(r_1, r_3)$ is $O(\ell^2)$ for $r_1, r_3 \gtrsim r_*$.

To conclude that $A_{S[K,K;0]} + A_{S[K,K;K]}$ is at most $O((\ell^2/r_*^2) \log(r_*/\ell))$, we have to examine whether $[\mathcal{O}_{r_1}^{[1]} \mathcal{O}_{r_3}^{[2]} v(r_1, r_3)]_{r_1=r_3=r}$ behaves well at $r \rightarrow \infty$ and at $r \rightarrow 0$ so that the operation of $\int d^3x' R^{-2} (R + \ell)$ is well-defined.

First we consider the large r limit. Since r_2 and r_4 are bounded by the radius of the star, $\mu(R_{12} + R_{34})$ can be replaced with $\mu(r_1 + r_3) \approx \ell/(r_1 + r_3)^2$ for large r_1 and/or r_3 , and hence $v(r_1, r_3) \approx \ell^2 (\kappa_4 M_*)^2 / r_1 r_3 (r_1 + r_3)^2$. The operation of $\mathcal{O}_{r_1}^{[1]}$ and $\mathcal{O}_{r_3}^{[2]}$ does not make the fall-off worse. For small r_1 ,

$$\begin{aligned} v(r_1, r_3) &= \frac{\kappa_4^2 \ell}{r_3} \int_0^{r_*} dr_2 r_2 \Sigma^{(1)}(r_2) \int_0^{r_*} dr_4 r_4 \Sigma^{(1)}(r_4) \int_{|r_3-r_4|}^{r_3+r_4} dR_{34} \left(\mu(r_2 + R_{34}) + \frac{r_1^2}{6} \mu''(r_2 + R_{34}) + O(r_1^4) \right) \\ &= \frac{\kappa_4^2 \ell}{r_3} \int_0^{r_*} dr_2 \left[\left(1 + \frac{r_1^2}{6} \partial_{r_2}^2 + O(r_1^4) \right) r_2 \Sigma^{(1)}(r_2) \right] \int_0^{r_*} dr_4 r_4 \Sigma^{(1)}(r_4) \int_{|r_3-r_4|}^{r_3+r_4} dR_{34} \mu(r_2 + R_{34}). \end{aligned} \quad (4.20)$$

Therefore, $\mathcal{O}_{r_1}^{[1]} v(r_1, r_3)$ is finite at $r_1 \rightarrow 0$.

For $A_{S[K,0;K]}$, a similar expression is obtained as

$$A_{S[K,0;K]} = \frac{2}{r_*^2} \int dr' r' \left[\mathcal{O}_{r_3}^{[1]} w(r_1, r_2, r_3) \right]_{r_1=r, r_2=r_3=r'} \mathcal{O}_{r'}^{[2]} \mathcal{A}_0^{(1)}(r'), \quad (4.21)$$

with

$$\begin{aligned} w(r_1, r_2, r_3) &\equiv r_2 \int d\Omega_2 \int_0^\infty dy' G_K(\mathbf{x}_1, 0; \mathbf{x}_2, y') \mathcal{A}_K^{(1)}(r_3, y') \\ &= \frac{\kappa_4 \ell}{4r_1 r_3} \int_0^{r_*} dr_4 r_4 \Sigma^{(1)}(r_4) \int_{|r_1-r_2|}^{r_1+r_2} dR_{12} \int_{|r_3-r_4|}^{r_3+r_4} dR_{34} \mu(R_{12} + R_{34}), \end{aligned} \quad (4.22)$$

and $\mathcal{A}_0^{(1)}$ represents the part of $\mathcal{A}^{(1)}$ propagated by G_0 . In the same way, we can show that $A_{S[K,0;K]}$ is at most $O((\ell^2/r_*^2) \log(r_*/\ell))$.

2. $A_{Q[K,K;0]}$ and $A_{Q[K,K;K]}$

$A_{Q[K,K;0]}$ and $A_{Q[K,K;K]}$ are also bounded as

$$\begin{aligned} &|A_{Q[K,K;0]} + A_{Q[K,K;K]}| \\ &< \frac{1}{2\pi\ell} \int d^3x' \frac{R+\ell}{R^2} \int_0^\infty dy a^4(y) Q_{yy}(\mathbf{x}, y) \\ &\leq \frac{1}{8\pi\ell} \int d^3x' \frac{R+\ell}{R^2} \left\{ \left| \mathcal{A}_{,y}^{(1)} \mathcal{A}_K^{(1)} + \mathcal{B}_{,y}^{(1)} \mathcal{B}_K^{(1)} + 2\mathcal{C}_{,y}^{(1)} \mathcal{C}_K^{(1)} \right|_{y=0} \right. \\ &\quad \left. + \left| \int_0^\infty dy \left((a^4 \mathcal{A}_{,y}^{(1)})_{,y} \mathcal{A}_K^{(1)} + (a^4 \mathcal{B}_{,y}^{(1)})_{,y} \mathcal{B}_K^{(1)} + 2(a^4 \mathcal{C}_{,y}^{(1)})_{,y} \mathcal{C}_K^{(1)} \right) \right| \right\}. \end{aligned} \quad (4.23)$$

The first term in the curly brackets is at most $O((\ell^3/r_*^4) \log(r_*/\ell))$. By using the relations obtained from Eq. (2.32),

$$\frac{1}{a^4} (a^4 \mathcal{A}_{,y}^{(1)})_{,y} = -a^{-2} \Delta \mathcal{A}^{(1)} + 2\kappa_5 \Sigma^{(1)} \delta(y), \quad (4.24)$$

$$\frac{1}{a^4} (a^4 \mathcal{B}_{,y}^{(1)})_{,y} = a^{-2} \frac{1}{r} \partial_r \mathcal{A}^{(1)} - \frac{2\kappa_5}{r^3} \int dr r^2 \Sigma^{(1)} \delta(y), \quad (4.25)$$

and $\mathcal{C}^{(1)} = -(\mathcal{A}^{(1)} + \mathcal{B}^{(1)})/2$, the second term is reduced to the form in (4.16). Therefore, the contribution to A_Q from this term is at most $O((\ell^2/r_*^2) \log(r_*/\ell))$ for the same reason.

V. SUMMARY

In this paper we developed second order perturbations in the RS single brane model, restricting the configuration to the static axisymmetric one. From the 5-dimensional Einstein equations, we derived the master equations for second order perturbations. At the level of linear perturbations, we can use the RS gauge, in which all perturbation equations are decoupled. Since the transverse-traceless condition cannot be imposed on the second order, the second order perturbative equations are inevitably coupled. As we have shown, however, the 4-dimensional spatial components of the second order metric function $\mathcal{B}^{(2)}$ and $\mathcal{C}^{(2)}$ turned out to be concisely represented by the temporal component $\mathcal{A}^{(2)}$ with the first order metric functions. Therefore, the problem was reduced to solving a single differential equation for $\mathcal{A}^{(2)}$. Once we solve for $\mathcal{A}^{(2)}$, the other metric functions $\mathcal{B}^{(2)}$ and $\mathcal{C}^{(2)}$ follow from it. Further, to discuss the metric induced on the 3-brane, we introduced the GN gauge in which a hypersurface with constant 5th-coordinate coincides with the location of the 3-brane. we gave the second order gauge transformations between the RS gauge and the GN gauge explicitly.

Based on this formulation, we first discussed the zero mode truncation for second order perturbations. It was shown that the metric induced on the 3-brane evaluated by using the approximation of the zero mode truncation exactly agrees with that for the 4-dimensional Einstein gravity.

Next, we evaluated the contribution to the metric functions from the KK modes. Since the mode-by-mode analysis shows a pathological feature even at the level of linear perturbations, it is necessary to sum up all the mass eigenvalues

to handle the KK mode interactions. Performing such an analysis, we have confirmed that the correction due to the KK mode coupling on the induced metric is suppressed by a factor of $O((\ell^2/r_*^2) \log(r_*/\ell))$, and there appears no pathological behavior. We therefore conclude that second order perturbations in the RS single brane model behave well and the result basically agrees with the prediction by the 4-dimensional Einstein gravity. The relative order of the correction is $O((\ell^2/r_*^2) \log(r_*/\ell))$. In the language of the post-Newtonian (PN) analysis of the 4-dimensional Einstein gravity, the order of $\mathcal{A}^{(2)}$ and $\mathcal{B}^{(1)}$ is 1PN, and that of $\mathcal{B}^{(2)}$ is 2PN. According to the parameterized post-Newtonian (PPN) formalism, the PPN parameters at the 1PN order, β and γ , are observationally constrained to the accuracy of about 0.1% or so [32]. Although this accuracy might be improved in the future, there will be no chance, unfortunately, to find the evidence for the large extra-dimension from the precision test of the solar system since the predicted deviation in β, γ is $O((\ell^2/r_*^2) \log(r_*/\ell))$.

Of course, since our discussion developed in the present paper is restricted to second order perturbations with spherical symmetry, the gravity induced on the 3-brane in the RS single brane model might deviate from the 4-dimensional Einstein gravity in more general situation. Hence, it would be necessary to develop a more complete proof of the coincidence including all higher order terms would. Although we did not discuss the RS two brane model, it is also interesting to study the post Newtonian correction in this model. The analysis is now in progress and will be reported soon in a separate paper.

Acknowledgements

H.K. would like to thank Takashi Nakamura and Hideo Kodama for informative comments and discussion. We also like to thank V.F. Mukhanov for informing us Ref. [34]. This work is supported by the Monbukagakusho Grant-in-Aid No. 1270154.

APPENDIX A: DERIVATION OF $\bar{\mathcal{B}}^{(2)}$

As was mentioned in the text below Eq. (4.4), the expression given in the last line of Eq. (4.4) contain the terms with y -integration. They are explicitly written as

$$\Delta^{-1} \left(-\frac{3}{\ell} \int_0^\infty dy a^2 Q_{yy} + \frac{1}{r^2} \partial_r \int_0^\infty dy r^2 Q_{yr} \right). \quad (\text{A1})$$

The first integral comes from $\xi_y^{(2)}$ contained in $\delta\mathcal{A}^{(2)}$, $\delta\mathcal{B}^{(2)}$ and $\delta\mathcal{C}^{(2)}$. In obtaining the expression (4.5) from (4.4), how to reduce these terms will be the only non-trivial manipulation. The rest of computation is slightly complicated but almost straightforward.

To rewrite the first integral, we use the relation

$$\begin{aligned} -\frac{1}{2} Q_{yy}(r, 0) &= \frac{1}{8} \int_0^\infty dy \partial_y \left\{ a^{-6} [(a^4 \mathcal{A}_{,y}^{(1)})^2 + (a^4 \mathcal{B}_{,y}^{(1)})^2 + 2(a^4 \mathcal{C}_{,y}^{(1)})^2] \right\}, \\ &= \frac{3}{\ell} \int_0^\infty dy a^2 Q_{yy} + \frac{1}{8} \int_0^\infty dy \frac{1}{a^2} [(3\mathcal{A}_{,y}^{(1)} + \mathcal{B}_{,y}^{(1)})(a^4 \mathcal{A}_{,y}^{(1)})_{,y} + (3\mathcal{B}_{,y}^{(1)} + \mathcal{A}_{,y}^{(1)})(a^4 \mathcal{B}_{,y}^{(1)})_{,y}], \end{aligned}$$

where we have used the traceless condition $\mathcal{C}^{(1)} = -(\mathcal{A}^{(1)} + \mathcal{B}^{(1)})/2$. On the other hand, the second part is rewritten as

$$\frac{1}{r^2} \partial_r \int_0^\infty dy r^2 Q_{yr} = -\frac{1}{2} Q_{rr} + \frac{1}{8} \int_0^\infty dy [(3\mathcal{A}_{,y}^{(1)} + \mathcal{B}_{,y}^{(1)}) \Delta \mathcal{A}^{(1)} + (3\mathcal{B}_{,y}^{(1)} + \mathcal{A}_{,y}^{(1)}) \Delta \mathcal{B}^{(1)}],$$

where again we have used the traceless condition. After the substitution of these relations and (4.24), the remaining y -integration is just

$$\frac{1}{8} \int_0^\infty dy (3\mathcal{B}_{,y}^{(1)} + \mathcal{A}_{,y}^{(1)}) \left[\Delta + \frac{1}{a^2} \partial_y a^4 \partial_y \right] \mathcal{B}^{(1)}. \quad (\text{A2})$$

We can derive

$$\frac{1}{a^2} (a^4 \mathcal{B}_{,y}^{(1)})_{,y} = - \left(\Delta \mathcal{B}^{(1)} + \frac{2}{r} \mathcal{B}_{,r}^{(1)} \right) - \frac{2\kappa_5}{r^3} \int dr r^2 \Sigma^{(1)} \delta(y), \quad (\text{A3})$$

from Eq. (4.25), and $(3\mathcal{B}_{,y}^{(1)} + \mathcal{A}_{,y}^{(1)}) = -r\mathcal{B}_{,r}^{(1)}$ from Eq. (2.18). With the aid of these relations, it is easy to see that the remaining y -integration (A2) can be performed.

APPENDIX B: SECOND ORDER PERTURBATIONS IN THE 4D EINSTEIN GRAVITY

In this Appendix, we give second order perturbations in the 4-dimensional Einstein gravity for the comparison with the zero mode truncation of the gravity in the RS single brane model. Although the results are well known as the 2nd post-Newtonian analysis [33], we present a brief derivation for the following two reasons. 1) Since we are working in a specific gauge, we need to consider a gauge transformation to compare our results with the expression presented in a different gauge. 2) To compute metric perturbations in our restricted situation from the beginning is much easier than to follow a literature in which unrestricted cases are discussed.

We assume that the 4-dimensional metric is static and isotropic,

$$ds^2 = -e^{A(r)}dt^2 + e^{B(r)}d\mathbf{x}^2.$$

Up to the second order, the 4-dimensional Einstein equations with the energy momentum tensor given in (2.2) are

$$R_t^t = \frac{1}{2}(B-1)\Delta A - \frac{1}{4}A_{,r}(A+B)_{,r} = -\frac{\kappa_4}{2}(\rho + 3P), \quad (\text{B1})$$

$$R_r^r = \frac{1}{2}(B-1)\left(\Delta(A+2B) - \frac{2}{r}(A+B)_{,r}\right) + \frac{1}{4}A_{,r}(B-A)_{,r} = \frac{\kappa_4}{2}(\rho - P), \quad (\text{B2})$$

$$R_\theta^\theta = \frac{1}{2}(B-1)\left(\Delta B + \frac{1}{r}(A+B)_{,r}\right) - \frac{1}{4}B_{,r}(A+B)_{,r} = \frac{\kappa_4}{2}(\rho - P). \quad (\text{B3})$$

We expand A and B to the second order as (2.9). Solving these equations at the linear order, we obtain $A^{(1)} = -B^{(1)} = 2\phi$, where $\phi(r)$ is similarly defined as in (3.10). Putting these results into Eq. (B1), the equation for $A^{(2)}$ becomes

$$\Delta A^{(2)} = \kappa_4(\rho^{(2)} + 3P) - 4\phi\Delta\phi, \quad (\text{B4})$$

which agrees with (4.8). Eliminating $\Delta A^{(2)}$ and P from Eqs. (B1), (B2) and (B3), we obtain

$$\Delta B^{(2)} = -\kappa_4\rho^{(2)} + 4\phi\Delta\phi - (\phi_{,r})^2. \quad (\text{B5})$$

Hence it was shown that the result obtained by the zero mode truncation (4.9) agrees with that for the 4-dimensional Einstein gravity (B5).

- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429**, 263 (1998); Phys. Rev. D **59**, 086004 (1999).
- [2] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436**, 257 (1998).
- [3] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
- [5] J. Garriga and T. Tanaka, Phys. Rev. Lett. **84**, 2778 (2000).
- [6] T. Shiromizu, K. Maeda and M. Sasaki, Phys. Rev. D **62**, 024012 (2000).
- [7] M. Sasaki, T. Shiromizu and K. Maeda, Phys. Rev. D **62**, 024008 (2000).
- [8] U. Gen and M. Sasaki, gr-qc/0011078.
- [9] T. Tanaka and X. Montes, Nucl. Phys. **B582**, 259 (2000).
- [10] S. B. Giddings, E. Katz and L. Randall, JHEP **0003**, 023 (2000).
- [11] T. Tanaka, Prog. Theor. Phys. **104**, 545 (2000).
- [12] C. Csaki, M. Graesser, L. Randall and J. Terning, Phys. Rev. D **62**, 045015 (2000).
- [13] C. Charmousis, R. Gregory and V. A. Rubakov, Phys. Rev. D **62**, 067505 (2000).
- [14] T. Chiba, Phys. Rev. D **62**, 021502 (2000).
- [15] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999).
- [16] W. D. Goldberger and M. B. Wise, Phys. Lett. **B475**, 275 (2000).
- [17] O. DeWolfe, D. Z. Freedman, S. S. Gubser and A. Karch, Phys. Rev. D **62**, 046008 (2000).
- [18] H. Kodama, A. Ishibashi and O. Seto, Phys. Rev. D **62**, 064022 (2000).
- [19] S. Mukohyama, Phys. Rev. D **62**, 084015 (2000)., Class. Quant. Grav. **17**, 4777 (2000).
- [20] R. Maartens, Phys. Rev. D **62**, 084023 (2000).
- [21] K. Koyama and J. Soda, Phys. Rev. D **62**, 123502 (2000).

- [22] C. van de Bruck, M. Dorca, R. H. Brandenberger and A. Lukas, Phys. Rev. D **62** 123515 (2000).
- [23] D. Langlois, Phys. Rev. D **62**, 126012 (2000).
- [24] A. Chamblin and G. W. Gibbons, Phys. Rev. Lett. **84**, 1090 (2000).
- [25] A. Chamblin, S. W. Hawking and H. S. Reall, Phys. Rev. D **61**, 065007 (2000).
- [26] R. Gregory and R. Laflamme, Phys. Rev. Lett. **70** 2837 (1993).
- [27] R. Gregory, Class. Quant. Grav. **17**, L125 (2000).
- [28] A. Chamblin, H. S. Reall, H. Shinkai and T. Shiromizu, Phys. Rev. D **63**, 064015 (2001).
- [29] R. Emparan, G. T. Horowitz and R. C. Myers, JHEP**0001**, 007 (2000).
- [30] N. Dadhich, R. Maartens, P. Papadopoulos and V. Rezanian, Phys. Lett. B **487**, 1 (2000).
- [31] W. Israel, Nuovo Cim. **44B**, 1 (1966).
- [32] C.M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, Cambridge 1993), revised ed; Lecture notes from the 1998 Slac Summer Institute on Particle Physics, gr-qc/9811036.
- [33] e.g., S. Chandrasekhar and Y. Nutku, Astroph. J. **158**, 55 (1969); L. Blanchet, Phys. Rev. D **51**, 2559 (1995).
- [34] I. Giannakis and H. Ren, Phys. Rev. D **63**, 024001 (2001)